arXiv:math/0501528v1 [math.NT] 28 Jan 2005

# TRANSFORMATIONS OF RAMANUJAN'S SUMMATION FORMULA AND ITS APPLICATIONS

Chandrashekar Adiga[1] and N.Anitha [2]
Department of Studies in Mathematics
University of Mysore, Manasagangotri
Mysore-570006, INDIA.

Taekyun Kim [3]
Institute of Science Education,
Kongju National University,
Kongju 314-701,
KOREA.

**Abstract:** In this paper, we obtain some new transformation formulas for Ramanujan's ${}_1\psi_1$ summation formula and also establish some eta-function identities. We also deduce a $q$- gamma function identity, an $q$-integral and some interesting series representations for $\frac{\pi^{3/2}}{2\sqrt{2}\Gamma^2(3/4)}$ and the beta function $B(x,y)$ .

**Keywords and Phrases:** Basic hypergeometric series, Dedekind eta-function, $q$-gamma function, $q$-integrals.

**2000 AMS Subject Classification:** 33D15, 11F20, 33D05.

# 1 Introduction :

One of the most celebrated identities of Ramanujan is his ${}_1\psi_1$-summation formula which can be stated as follows:

$$\sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n = \frac{(az)_\infty (q/az)_\infty (q)_\infty (b/a)_\infty}{(z)_\infty (b/az)_\infty (b)_\infty (q/a)_\infty} \tag{1.1}$$

where $\mid b/a \mid < \mid z \mid < 1$. This identity was first brought before the mathematical world by G. H. Hardy [7] who described it as "A remarkable formula with many parameters". Hardy did not supply a proof but indicated that a proof could be constructed from

---

[1] e-mail : c_adiga@hotmail.com
[2] e-mail : ani_nie@indiatimes.com
[3] e-mail : tkim@kongju.ac.kr

the $q$-binomial theorem. There are several proofs of (1.1) in the literature [c.f 1 ]. Most proofs of (1.1) found in the literature do depend on the $q$-binomial theorem. The transformations of the basic hypergeometric series are extremely useful in the theory of partitions. The main purpose of this paper is to obtain some transformations for Ramanujan's ${}_1\psi_1$ summation formula (1.1).

As customary we employ the following definitions and notations:

$$(a)_\infty := (a;q)_\infty = \prod_{n=0}^{\infty}(1-aq^n), \quad \mid q \mid < 1, \tag{1.2}$$

$$(a)_n := \frac{(a)_\infty}{(aq^n)_\infty}, \quad n : \text{any integer}. \tag{1.3}$$

The generalized basic hypergeometric series is defined by

$${}_r\phi_s\left[\begin{array}{l} a_1, a_2, ..., a_r; q; z \\ b_1, b_2, ..., b_s \end{array}\right] = \sum_{n=0}^{\infty}\frac{(a_1)_n(a_2)_n...(a_r)_n}{(q)_n(b_1)_n(b_2)_n...(b_s)_n}z^n, \quad \mid z \mid < 1, \quad \mid q \mid < 1. \tag{1.4}$$

The bilateral basic hypergeometric series is defined by

$${}_r\psi_r\left[\begin{array}{l} a_1, a_2, ..., a_r; q; z \\ b_1, b_2, ..., b_r \end{array}\right] = \sum_{n=-\infty}^{\infty}\frac{(a_1)_n(a_2)_n...(a_r)_n}{(b_1)_n(b_2)_n...(b_r)_n}z^n, \tag{1.5}$$

where $\left|\frac{b_1...b_r}{a_1...a_r}\right| < \mid z \mid < 1, \mid q \mid < 1.$

In Section 3 we obtain some special cases of the transformations for Ramanujan's ${}_1\psi_1$ summation formula obtained in Section 2 and in Section 4 we also deduce some eta-function identities. In the final section we derive some interesting series representations for the beta function $B(x, y)$ and $\frac{\pi^{3/2}}{2\sqrt{2}\Gamma^2(3/4)}$.

## 2 Main Results

To prove our main result we need the Heine's transformation[5, Eq.(III.1) p.241, Eq.(III.2), p.241]

$$\sum_{n=0}^{\infty}\frac{(a)_n(b)_n}{(q)_n(c)_n}z^n = \frac{(b)_\infty(az)_\infty}{(c)_\infty(z)_\infty}\sum_{n=0}^{\infty}\frac{(c/b)_n(z)_n}{(q)_n(az)_n}b^n \tag{2.1}$$

$$= \frac{(c/b)_\infty(bz)_\infty}{(c)_\infty(z)_\infty}\sum_{n=0}^{\infty}\frac{(abz/c)_n(b)_n}{(q)_n(bz)_n}(c/b)^n. \tag{2.2}$$

**Theorem 2.1.** If $\mid b/a \mid < \mid z \mid < 1$ and $\mid q \mid < \min \{1, \quad \mid b \mid\}$ then,

$$\sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n = -1 + \frac{(b/a)_\infty (az)_\infty}{(b)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(a)_n (aqz/b)_n}{(q)_n (az)_n} (b/a)^n$$
$$+ \frac{(q/b)_\infty (bq/az)_\infty}{(q/a)_\infty (b/az)_\infty} \sum_{n=0}^{\infty} \frac{(b/a)_n (b/az)_n}{(q)_n (bq/az)_n} (q/b)^n. \tag{2.3}$$

**Proof:** We have

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n &= \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} z^n + \sum_{n=1}^{\infty} \frac{(q/b)_n}{(q/a)_n} (b/az)^n \\ &= -1 + \sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} z^n + \sum_{n=0}^{\infty} \frac{(q/b)_n}{(q/a)_n} (b/az)^n. \end{aligned} \tag{2.4}$$

Putting $a = q$, $b = a$ and $c = b$ in (2.2) we obtain

$$\sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} z^n = \frac{(b/a)_\infty (az)_\infty}{(b)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(a)_n (aqz/b)_n}{(q)_n (az)_n} (b/a)^n. \tag{2.5}$$

Changing $a$ to $q$, $b$ to $q/b$, $c$ to $q/a$ and $z$ to $b/az$ in (2.1) we obtain

$$\sum_{n=0}^{\infty} \frac{(q/b)_n}{(q/a)_n} (b/az)^n = \frac{(q/b)_\infty (bq/az)_\infty}{(q/a)_\infty (b/az)_\infty} \sum_{n=0}^{\infty} \frac{(b/a)_n (b/az)_n}{(q)_n (bq/az)_n} (q/b)^n. \tag{2.6}$$

Substituting (2.5) and (2.6) in (2.4) we obtain (2.3).

**Theorem 2.2.** If $\mid \frac{b}{a} \mid < \mid z \mid < 1$ and $\mid a \mid < 1$ then,

$$\sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n = -1 + \frac{(a)_\infty (qz)_\infty}{(b)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(b/a)_n (z)_n}{(q)_n (qz)_n} (a)^n$$
$$+ \frac{(b/a)_\infty (q/az)_\infty}{(q/a)_\infty (b/az)_\infty} \sum_{n=0}^{\infty} \frac{(q/z)_n (q/b)_n}{(q)_n (q/az)_n} (b/a)^n. \tag{2.7}$$

**Proof:** Changing $a$ to $q$, $b$ to $a$ and then $c$ to $b$ in (2.1), we deduce

$$\sum_{n=0}^{\infty} \frac{(a)_n}{(b)_n} z^n = \frac{(a)_\infty (qz)_\infty}{(b)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(b/a)_n (z)_n}{(q)_n (qz)_n} (a)^n. \tag{2.8}$$

Putting $a = q$, $b = q/b$, $c = q/a$ and $z = b/az$ in (2.2), we obtain

$$\sum_{n=0}^{\infty} \frac{(q/b)_n}{(q/a)_n} (b/az)^n = \frac{(b/a)_\infty (q/az)_\infty}{(q/a)_\infty (b/az)_\infty} \sum_{n=0}^{\infty} \frac{(q/z)_n (q/b)_n}{(q)_n (q/az)_n} (b/a)^n. \tag{2.9}$$

On employing (2.8) and (2.9) in (2.4) we obtain Theorem 2.2.

**Theorem 2.3.** If $\mid \frac{b}{a} \mid < \mid z \mid < 1$ then,

$$\sum_{n=-\infty}^{\infty} \frac{(a)_n}{(b)_n} z^n = -1 + \frac{(b/a)_\infty (az)_\infty}{(b)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(aqz/b)_n (a)_n}{(q)_n (az)_n} (b/a)^n + \frac{(b/a)_\infty (q/az)_\infty}{(q/a)_\infty (b/az)_\infty} \sum_{n=0}^{\infty} \frac{(q/z)_n (q/b)_n}{(q)_n (q/az)_n} (b/a)^n. \tag{2.10}$$

**Proof:** On employing (2.5) and (2.9) in (2.4) we obtain Theorem 2.3.

# 3 Special Cases

Putting $a = -1/q$ and $b = -1$ in (2.3), we obtain

$$\sum_{n=-\infty}^{\infty} \frac{z^n}{1+q^{n-1}} = \frac{-q}{1+q} + \frac{q}{1+q} \frac{(q)_\infty (-z/q)_\infty}{(-1)_\infty (z)_\infty} \sum_{n=0}^{\infty} \frac{(z)_n (-1/q)_n}{(q)_n (-z/q)_n} q^n + q \sum_{n=0}^{\infty} \frac{1}{1 - \frac{q^{n+1}}{z}} (-q)^n, \tag{3.1}$$

where $\mid q \mid < \mid z \mid < 1$ .

Changing $b$ to $-q^3$, $a$ to $-q$, $z$ to $q$ in (2.7) we get,

$$\sum_{n=-\infty}^{\infty} \frac{q^n}{(1+q^{n+1})(1+q^{n+2})} = \frac{(1+q^2)(1+q)}{q(1-q)}. \tag{3.2}$$

Putting $b = -q^2$, $a = -1/q$ and $z = q^{1/2}$ in (2.10) and then changing $q$ to $q^2$, we obtain

$$\sum_{n=-\infty}^{\infty} \frac{2(1+1/q^2)(1+q^2)q^n}{(1+q^{2n-2})(1+q^{2n})(1+q^{2n+2})} = -1 + \frac{(q^6;q^2)_\infty (-1/q;q^2)_\infty}{(-q^4;q^2)_\infty (q;q^2)_\infty} \sum_{n=0}^{\infty} \frac{(1/q^3;q^2)_n (-1/q^2;q^2)_n}{(q^2;q^2)_n (-1/q;q^2)_n} q^{6n} + \frac{(q^6;q^2)_\infty (-q^3;q^2)_\infty}{(-q^4;q^2)_\infty (q^5;q^2)_\infty} \sum_{n=0}^{\infty} \frac{(q;q^2)_n (-1/q^2;q^2)_n}{(q^2;q^2)_n (-q^3;q^2)_n} q^{6n}. \tag{3.3}$$

# 4 Eta-function identities

The Dedekind eta-function is defined by

$$\begin{aligned}\eta(\tau) &:= e^{\pi i\tau/12}\prod_{n=1}^{\infty}(1-e^{2\pi in\tau})\\ &= q^{1/24}(q;q)_\infty, \qquad (4.1)\end{aligned}$$

where $q = e^{2\pi i\tau}$ and Im $\tau > 0$.

The eta-function is useful in the study of modular forms. Berndt and L. C. Zhang [3] have obtained a number of eta-function identities found in Ramanujan's notebook employing the theory of modular forms. In [4], Bhargava and Somashekara show how a family of interesting eta-function identities can be obtained from a limiting case of the Ramanujan's ${}_1\psi_1$-summation formula.
Now we derive some new $\eta$- function identities.

Employing Ramanujan's ${}_1\psi_1$ summation formula in (2.3) and then putting $a = 1/q$, $b = q^{1/2}$, $z = q^{1/2}$ and changing $q$ to $q^2$, we deduce

$$\frac{\eta(\tau)}{\eta^2(2\tau)} = q^{-1/8} - \frac{q^{7/8}}{1+q}\frac{(q;q^2)_\infty}{(q^2;q^2)_\infty}\sum_{n=0}^{\infty}\frac{(q^3;q^2)_n}{(q^4;q^2)_n}q^n. \qquad (4.2)$$

This can also be obtained from $q$-binomial theorem.

Employing Ramanujan's ${}_1\psi_1$ summation formula in (2.7) and then putting $a = -q$, $b = -q^3$, $z = q^{1/2}$ and changing $q$ to $q^2$, we deduce

$$\frac{\eta^{10}(2\tau)}{\eta^4(\tau)\eta^2(4\tau)} = \frac{2(1+q)q^{4/3}}{(1+q^2)(1+q^4)} - \frac{2q^{4/3}}{1-q}\sum_{n=0}^{\infty}\frac{1-q^{2n+2}}{1-q^{2n+1}}(-q^2)^n - \\ \frac{2(1+q)q^{4/3}}{(1+q^2)(1+q^4)}\frac{(q^4;q^2)_\infty(-1/q;q^2)_\infty}{(-1;q^2)_\infty(q^3;q^2)_\infty}\sum_{n=0}^{\infty}\frac{(q;q^2)_n(-1/q^4;q^2)_n}{(q^2;q^2)_n(-1/q;q^2)_n}q^{4n}. \qquad (4.3)$$

Employing Ramanujan's ${}_1\psi_1$ summation formula in (2.3) and then putting $a = -q^{-5/3}$, $b = -q^{1/3}$, $z = q^{4/3}$ and changing $q$ to $q^3$, we deduce

$$\frac{\eta^3(3\tau)}{\eta(\tau)} = \frac{q^{4/3}(1+q+q^2)}{(1+q^2)(1+q^5)}\left[-1+\frac{(q^6;q^3)_\infty(-1/q;q^3)_\infty}{(-q;q^3)_\infty(q^4;q^3)_\infty}\sum_{n=0}^{\infty}\frac{(q;q^3)_n(-1/q^5;q^3)_n}{(q^3;q^3)_n(-1/q;q^3)_n}q^{6n}\right]$$
$$+\frac{q^{4/3}(1+q+q^2)}{(1+q^2)(1+q^5)}\frac{(q^6;q^3)_\infty(-q^4;q^3)_\infty}{(-q^8;q^3)_\infty(q^2;q^3)_\infty}\sum_{n=0}^{\infty}\frac{(1/q;q^3)_n(-q^2;q^3)_n}{(q^3;q^3)_n(-q^4;q^3)_n}q^{6n}. \tag{4.4}$$

# 5 Some q-Gamma Function Identities:

F.H. Jackson [8] defined the $q$-analogue of the gamma function by

$$\Gamma_q(x) = \frac{(q)_\infty}{(q^x)_\infty}(1-q)^{1-x}, \quad 0<q<1. \tag{5.1}$$

Jackson [8] also defined a $q$-integral by,

$$\int_0^a f(t)d_q(t) = a(1-q)\sum_{n=0}^{\infty} f(aq^n)q^n \tag{5.2}$$

and

$$\int_0^\infty f(t)d_q(t) = (1-q)\sum_{n=-\infty}^{\infty} f(q^n).q^n. \tag{5.3}$$

In this section, as an application of the transformation formulas of Section 2, we deduce some $q$-gamma function identities and an $q$-integral which give interesting series representations for the beta function B(x,y) and $\frac{\pi^{3/2}}{2\sqrt{2}\Gamma^2(3/4)}$.

**Theorem 5.1.** If $0<z<b-a<1$ and $b<1$ then,

$$\frac{\Gamma_q(b)\Gamma_q(1-a)\Gamma_q(z)\Gamma_q(b-a-z)}{\Gamma_q(b-a)\Gamma_q(a+z)\Gamma_q(1-a-z)} = -(1-q)^{a+1-b}+(1-q)^{a+1-b}\frac{\Gamma_q(b)\Gamma_q(z)}{\Gamma_q(b-a)\Gamma_q(a+z)}$$
$$\sum_{n=0}^{\infty}\frac{(q^{a+1+z-b})_n(q^a)_n}{(q)_n(q^{a+z})_n}(q^{b-a})^n+\frac{\Gamma_q(1-a)\Gamma_q(b-a-z)}{\Gamma_q(1-b)\Gamma_q(b+1-a-z)}\sum_{n=0}^{\infty}\frac{(q^{b-a})_n(q^{b-z-a})_n}{(q^{b+1-a-z})_n(q)_n}(q^{1-b})^n. \tag{5.4}$$

**Proof:** Changing $a$ to $q^a$, $b$ to $q^b$ and $z$ to $q^z$ in (2.3) we obtain the required result.

**Corollary:** In Theorem 5.1 if we let $q \to 1$ and put $a=0$ we deduce,

$$\frac{\Gamma(1-b)\Gamma(b+1-z)}{\Gamma(1-z)} = \sum_{n=0}^{\infty}\frac{(b)_n}{n!}\frac{b-z}{b-z+n}. \tag{5.5}$$

**Remark:** Putting $b - z = y$ and $1 - b = x$ in (5.5) we obtain a well known [6] representation for the beta function $B(x, y)$ given by,

$$B(x,y) = \sum_{n=0}^{\infty} \frac{\prod_{k=1}^{n}(k-x)}{n!} \frac{1}{n+y}. \tag{5.6}$$

**Theorem 5.2.** If $0 < z < b - a < 1$ and $a > 0$ then,

$$\frac{\Gamma(a)\Gamma(z)\Gamma(1-a)\Gamma(b-a-z)}{\Gamma(a+z)\Gamma(b-a)\Gamma(1-a-z)} = \sum_{n=0}^{\infty} \frac{(b-a)_n}{n!(n+z)}. \tag{5.7}$$

**Proof:** Changing $a$ to $q^a$, $b$ to $q^b$ and $z$ to $q^z$ in (2.7) we obtain

$$\frac{\Gamma_q(z)\Gamma_q(b-a-z)\Gamma_q(b)\Gamma_q(1-a)}{\Gamma_q(a+z)\Gamma_q(1-a-z)\Gamma_q(b-a)} = -(1-q)^{a+1-b} + \frac{\Gamma_q(b)\Gamma_q(z)}{\Gamma_q(a)\Gamma_q(z+1)}$$
$$\sum_{n=0}^{\infty} \frac{(q^{b-a})_n (z)_n}{(q)_n (q^{1+z})_n} (q^a)^n + (1-q)^{a+1-b} \frac{\Gamma_q(1-a)\Gamma_q(b-a-z)}{\Gamma_q(1-a-z)\Gamma_q(b-a)} \sum_{n=0}^{\infty} \frac{(q^{1-z})_n (q^{1-b})_n}{(q)_n (q^{1-a-z})_n} (q^{b-a})^n. \tag{5.8}$$

Letting $q \to 1$ in (5.8) we obtain Theorem 5.2.

**Remark:** Putting $a = 1/4$, $b = 1$ and $z = 1/2$ in Theorem 5.2 we obtain,

$$\frac{\pi^{3/2}}{2\sqrt{2}\Gamma^2(3/4)} = \sum_{n=0}^{\infty} \frac{(1/2)_n}{n!(4n+1)} \tag{5.9}$$

which is very similar to the result obtained by Ramanujan[2, p.24].

**Theorem 5.3.** If $0 < z < b - a < 1$ then,

$$\begin{aligned} \frac{\Gamma_q(b)\Gamma_q(1-a)\Gamma_q(z)\Gamma_q(b-a-z)}{\Gamma_q(b-a)\Gamma_q(a+z)\Gamma_q(1-a-z)} &= -(1-q)^{a+1-b} + \frac{\Gamma_q(1-a)\Gamma_q(b-a-z)}{\Gamma_q(b-a)\Gamma_q(1-z)\Gamma_q(1-b)} \\ \int_0^1 f(x)d_q(x) &+ \frac{\Gamma_q(b)\Gamma_q(z)}{\Gamma_q(b-a)\Gamma_q(a+1+z-b)\Gamma_q(a)} \int_0^1 g(x)d_q(x) \end{aligned} \tag{5.10}$$

where,

$$f(x) = \frac{(xq)_\infty (xq^{1-a-z})_\infty}{(xq^{1-z})_\infty (xq^{1-b})_\infty} x^{b-a-1}$$

and

$$g(x) = \frac{(xq)_\infty (xq^{a+z})_\infty}{(xq^{a+1+z-b})_\infty (xq^a)_\infty} x^{b-a-1}.$$

**Proof:** Changing $a$ to $q^a$, $b$ to $q^b$ and $z$ to $q^z$ in (2.10) we get,

$$\frac{\Gamma_q(b)\Gamma_q(1-a)\Gamma_q(z)\Gamma_q(b-a-z)}{\Gamma_q(b-a)\Gamma_q(a+z)\Gamma_q(1-a-z)}(1-q)^{b-a-1} = -1 + \frac{\Gamma_q(1-a)\Gamma_q(b-a-z)}{\Gamma_q(b-a)\Gamma_q(1-z)\Gamma_q(1-b)}$$
$$(1-q)^{(b-a)}\sum_{n=0}^{\infty}\frac{(q^{n+1})_\infty(q^{n+1-a-z})_\infty}{(q^{n+1-z})_\infty(q^{n+1-b})_\infty}(q^{b-a})^n + \frac{\Gamma_q(b)\Gamma_q(z)}{\Gamma_q(b-a)\Gamma_q(a+1+z-b)\Gamma_q(a)}$$
$$(1-q)^{(b-a)}\sum_{n=0}^{\infty}\frac{(q^{n+1})_\infty(q^{n+a+z})_\infty}{(q^{n+a+1+z-b})_\infty(q^{n+a})_\infty}(q^{b-a})^n \tag{5.11}$$

Applying (5.2) to the right side of (5.11) we obtain Theorem 5.3.
**Corollary:** In Theorem 5.3 if we let $q \to 1$ we deduce,

$$\frac{\Gamma(b)\Gamma(1-a)\Gamma(z)\Gamma(b-a-z)}{\Gamma(a+z)\Gamma(1-a-z)} =$$
$$\left[\frac{\Gamma(1-a)\Gamma(b-a-z)}{\Gamma(1-z)\Gamma(1-b)} + \frac{\Gamma(b)\Gamma(z)}{\Gamma(a)\Gamma(a+1+z-b)}\right] B(b-a, a-b+1). \tag{5.12}$$

# References

[1] C. Adiga, B. C. Berndt, S. Bhargava and G. N. Watson, Chapter 16 of Ramanujan's second notebook: *Theta-functions and q-series*, Mem.Amer.Math.Soc.No.315, 53(1985), 1-85.

[2] B. C. Berndt, *Ramanujan's Notebooks II*, Springer-Verlag, New York, 1991.

[3] B. C. Berndt and L. C. Zhang, *Ramanujan's identities for eta functions*, Math.Ann.292(3) (1992), 561-573.

[4] S. Bhargava and D. D. Somashekara, *Some eta function identities deducible from Ramanujan's* ${}_1\psi_1$*Summation*, J.Math.Anal.Appl.176(2)(1993), 554-560.

[5] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press, 1990.

[6] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals*, Series and Products, 4th ed., Academic Press, New York, 1965.

[7] G. H. Hardy, *Ramanujan*, 3rd ed. Chelsea, NewYork, 1978.

[8] F. H. Jackson, *On q-definite integrals*, Quar.Jour.Pure and Applied Math. 41(1910), 193-203.